arXiv:math/0508304v1 [math.AC] 17 Aug 2005

On the equations defining curves in a polynomial algebra

Ze Min Zeng
Department of Mathematics,
Washington University in St. Louis,
St. Louis, Missouri, 63130
Email: `zmzeng@math.wustl.edu`

Abstract

Let A be a commutative Noetherian ring of dimension n ($n \geq 3$). Let I be a local complete intersection ideal in $A[T]$ of height n. Suppose I/I^2 is free $A[T]/I$-module of rank n and $(A[T]/I)$ is torsion in $K_0(A[T])$. It is proved in this paper that I is a set theoretic complete intersection ideal in $A[T]$ if one of the following conditions holds: (1) $n \geq 5$, odd; (2) n is even, and A contains the field of rational numbers; (3) $n = 3$, and A contains the field of rational numbers.

MSC: Primary 13C10; Secondary 13C40
Keywords: Set theoretic complete intersection, Euler class group

1 Introduction

Let A be a commutative Noetherian ring of dimension n. Let I be a local complete intersection ideal in $A[T]$ of height n. In [8], Mandal proved the following interesting theorem:

Theorem 1.1. *Let A be a commutative Noetherian ring of dimension n. Let I be a local complete intersection ideal in $A[T]$ of height n. Suppose I contains a monic polynomial. Then I is a set theoretic complete intersection in $A[T]$.*

Let A be a commutative Noetherian ring of dimension n. Let J be a local complete intersection ideal in $A[T]$ of height n. By the well known Ferrand-Szpiro construction[14], there exists a local complete intersection ideal I which is contained in J, such that $\sqrt{I} = \sqrt{J}$ and I/I^2 is free $A[T]/I$-module of rank n. So to show J is a set theoretic complete intersection in $A[T]$, it suffices to show I is a set theoretic complete intersection in $A[T]$. If A contains the field of rational numbers, in this paper, we are able to find a new sufficient condition for I to be a set theoretic complete intersection, which can be read as follows:

Theorem 1.2. *Let A be a commutative Noetherian ring of dimension n ($n \geq 3$) containing the field of rational numbers. Let I be a local complete intersection ideal in $A[T]$ of height n. Suppose I/I^2 is free $A[T]/I$-module of rank n and $(A[T]/I)$ is torsion in $K_0(A[T])$. Then I is a set theoretic complete intersection ideal in $A[T]$.*

If the dimension of A is odd and bigger than 5, the above theorem also holds without the condition A contains the field of rational numbers.

All rings in this paper are assumed to be commutative and Noetherian. All modules considered are assumed to be finitely generated.

2 Some preliminary results

In this section, we collect some well known results that will be used in the third section.

First let us begin with a standard definition.

Definition 2.1. Let A be a commutative Noetherian ring. $K_0(A)$ is called the *Grothendieck Group* of A, which is defined by taking the free abelian group generated by all finitely generated A-modules with finite projective dimension modulo the relation generated by $(M_2) = (M_1) + (M_3)$, whenever we have an exact sequence of finitely generated A-modules of finite projective dimension,

$$0 \to M_1 \to M_2 \to M_3 \to 0.$$

The following theorem is due to Boratynski[5] and Murthy[10]:

Theorem 2.2. *Let A be a commutative Noetherian ring of dimension n, and $I \subseteq A$ be a local complete intersection of height r ($r \leq n$). Suppose I/I^2 is A/I-free with base $\bar{f}_1, \ldots, \bar{f}_r$, $f_i \in I$, $\bar{f}_i$ is the class of f_i in I/I^2. Let $J = I^{(r-1)!} + (f_1, \ldots, f_{r-1})$. Then there exists a surjection $P \to J$ with P a projective A-module of rank r, such that $(P) - (A^r) = -(A/I)$ in $K_0(A)$.*

The next useful lemma which follows is due to Mohan Kumar[9, Lemma 1].

Lemma 2.3. *Let A be a ring and $J \subset A$ a finitely generated ideal. Suppose that J/J^2 is generated by n elements. Then for any $a \in A$, the ideal (J, a) is generated by $n+1$ elements.*

Next we state the cancellation theorem of Plumstead[11, Theorem 1]:

Theorem 2.4. *Let A be a commutative Noetherian ring of dimension n, let P and P' be projective $A[T]$-modules with rank $\geq n+1$. If $P \oplus A[T] \approx P' \oplus A[T]$, then $P \approx P'$.*

The following theorem is a result of Bass[1].

Theorem 2.5. *Let B be a ring with dimension $n+1$, P a stably free projective B-module of rank n, such that $P \oplus B \approx B^{n+1}$. If n is odd, then P has a unimodular element.*

The following theorem is a consequence of a result of Rao[13, Corollary 2.5] and Quillen's local-global principle[12, Theorem 1].

Theorem 2.6. *Let A be a Noetherian ring of dimension n. Suppose $n!$ is invertible in A. Then any projective module given by a unimodular row over $A[T]$ of length $n+1$ is extended from A. In other words, all stably free $A[T]$-module of rank n are extended from A.*

Let A be a commutative Noetherian ring of dimension n which contains the field of rational numbers. The Euler class group $E(A)$ of A is defined by Bhatwadekar and Sridharan in [3]. Many important and interesting theorems are proved in their paper. Let us quote one of them [3, Theorem 4.2]:

Theorem 2.7. *Let A be a Noetherian ring of dimension $n \geq 2$ which contains the field of rational numbers. Let J be an ideal of height n such that J/J^2 is generated by n elements, and let $\omega_J : (A/J)^n \to J/J^2$ be a local orientation of J. Suppose that the image of (J, ω_J) is zero in the Euler class group $E(A)$ of A. Then ω_J is a global orientation of J. In other words, ω_J can be lifted to a surjection $\psi : A^n \to J$.*

The following theorem is due to Das[6, Theorem 3.10]

Theorem 2.8. *Let A be a Noetherian ring of dimension $n \geq 3$, containing the field of rational numbers. Let $I \subset A[T]$ be an ideal of height n. Suppose there exists a surjection $\phi : A[T]^n \to I/I^2T$, such that $\phi \otimes A(T)$ can be lifted to a surjection $\phi' : A(T)^n \to IA(T)$. Then there is a surjection $\psi : A[T]^n \to I$ which lifts ϕ.*

3 Set theoretic complete intersection for curves in $A[T]$

Let A be a commutative Noetherian ring of dimension n. Let I be a local complete intersection ideal of height n in $A[T]$. Assume I/I^2 is free $A[T]/I$-module of rank n. If $(A[T]/I)$ is torsion in $K_0(A[T])$, then by the standard "thickening" method, we will show that we can find a local complete intersection ideal K in $A[T]$, such that $\sqrt{K} = \sqrt{I}$ and K is the surjective image of a stably free projective $A[T]$-module $\tilde{P}$ of rank n. More precisely:

Lemma 3.1. *Let A be a commutative Noetherian ring of dimension n ($n \geq 2$). Let I be a local complete intersection ideal of height n in $A[T]$, such that I/I^2 is free $A[T]/I$-module of rank n. If $(A[T]/I)$ is torsion in $K_0(A[T])$, then there exist a local complete intersection ideal K in $A[T]$, and a surjection $\xi : \tilde{P} \to K$ such that:*

1. *$K \subset I$, $\sqrt{K} = \sqrt{I}$,*
2. *$\tilde{P} \oplus A[T] \approx A[T]^{n+1}$,*
3. *$(A[T]/K) = 0$ in $K_0(A[T])$ and K/K^2 is free $A[T]/K$-module of rank n.*

Proof. We can find a regular sequence $f_1, \ldots, f_n$ in I, such that $I = (f_1, \ldots, f_n) + I^2$. Suppose $r(A[T]/I) = 0$ in $K_0(A[T])$. Let $J = I^r + (f_1, \ldots, f_{n-1})$, then $(A[T]/J) = r(A[T]/I) = 0$ in $K_0(A[T])$ and $J = (f_1, \ldots, f_{n-1}, f_n^r) + J^2$. Let $K = J^{(n-1)!} + (f_1, \ldots, f_{n-1})$, then K is a locally complete intersection ideal in $A[T]$, satisfying $(A[T]/K) = 0$ in $K_0(A[T])$, $\sqrt{I} = \sqrt{K}$ and K/K^2 is free $A[T]/K$-module of rank n generated by the image of $f_1, \ldots, f_{n-1}, f_n^{r(n-1)!}$. By theorem 2.2, there exists a projective $A[T]$-module $\tilde{P}$ of rank n and a surjection: $\tilde{P} \to K$, such that $(\tilde{P}) - (A[T]^n) = 0$ in $K_0(A[T])$. Hence $\tilde{P}$ is a stably free $A[T]$-module of rank n. By Plumstead cancellation theorem, $\tilde{P} \oplus A[T] \approx A[T]^{n+1}$. The proof of the lemma is complete.

□

Theorem 3.2. *Let A be a commutative Noetherian ring of dimension n ($n \geq 4$). Let I be a local complete intersection ideal of height n in $A[T]$, such that I/I^2 is free $A[T]/I$-module of rank n. Suppose $(A[T]/I)$ is torsion in $K_0(A[T])$ and n is odd. Then I is a set theoretic complete intersection in $A[T]$.*

Proof. By Lemma 3.1, there exist a local complete intersection ideal K in $A[T]$, and a surjection $\xi : \tilde{P} \to K$, such that $\sqrt{K} = \sqrt{I}$ and $\tilde{P}$ is a stably free projective $A[T]$-module of rank n, and $\tilde{P} \oplus A[T] \approx A[T]^{n+1}$. Since n is odd, by theorem 2.5, $\tilde{P}$ has a unimodular element. So we can write $\tilde{P} = A[T] \oplus \tilde{Q}$ where $\tilde{Q}$ is a stably free $A[T]$-module of rank $n-1$. Let $K_1 = \xi(\tilde{Q})$. By some suitable elementary transformation on $\tilde{P}$, we may assume $\mathrm{ht}K_1 = n-1$. Let $\xi((1,0)) = x$, then $(x, K_1) = K$. Since $n \geq 4$, by Bass cancellation theorem it is easy to see that $\tilde{Q}/K_1\tilde{Q}$ is free $A[T]/K_1$-module of rank $n-1$. Therefore, by lemma 2.3, $K = (K_1, x)$ is n generated, and hence K is a complete intersection. Thus I is a set theoretic complete intersection in $A[T]$. □

We need the following lemma to prove our next theorem.

Lemma 3.3. *Let B be a commutative Noetherian ring of dimension $n+1$ containing a field k. Let I be an ideal of height n which is a local complete intersection in B, such that I/I^2 is free B/I-module of rank n. Then there exists a regular sequence $f_1, \ldots, f_n$ in B and $s_1 \in I^2$ such that*

1. *$I = (f_1, \ldots, f_n, s_1)$, $s_1(1-s_1) \in (f_1, \ldots, f_n)$, $I = (f_1, \ldots, f_n) + I^2$, and*
2. *$\{f_1, \ldots, f_{n-1}, f_n - s_1^2\}$ is a regular sequence in B.*

Proof. There exists a regular sequence $f_1, \ldots, f_n$ in B such that $I = (f_1, \ldots, f_n) + I^2$. By Nakayama's lemma, there exists $s \in I$ such that $s(1-s) \in (f_1, \ldots, f_n)$ and $I = (f_1, \ldots, f_n, s)$. Since $s(1-s) \in (f_1, \ldots, f_n)$, we may further assume that $s \in I^2$. Notice that we can change s by $\prod_{i=1}^{m}(s - b_i f_n)$ for any positive integer m and $b_i \in B$. If $p_1, \ldots, p_t$ are the maximal elements in $\mathrm{Ass}(B/(f_1, \ldots, f_{n-1}))$, then $f_n \notin p_1, \ldots, p_t$.

If $s \in p_1, \ldots, p_t$, then $f_n - s^2 \notin p_1, \ldots, p_t$, and we are through.

If, say for example, $s \notin p_1$, but $s + bf_n \in p_1$ for some $b \in I$, we replace s by $s(s+bf_n)$ and assume $s \in p_1$. Repeating this procedure (that is, replacing s by $\prod_{i=1}^{m}(s - b_i f_n)$) and reordering p_i where $i \in \{1, \ldots, t\}$ if necessary, we may assume that $s \in p_1, \ldots, p_r$, $s - bf_n \notin p_k$ for $k > r$ and any $b \in I$.

Since $s \in p_1, \ldots, p_r$, $f_n - s^2 \notin p_1, \ldots, p_r$. If $f_n - s^2 \notin p_{r+1}, \ldots, p_t$, then we are done. So by reordering $p_{r+1}, \ldots, p_t$, we may assume $f_n - s^2 \notin p_{r+1}, \ldots, p_{r+l}$ and $f_n - s^2 \in p_{r+l+1}, \ldots, p_t$. Let $\lambda \in I \cap (\cap_{i=1}^{r+l} p_i) \setminus \cup_{j=r+l+1}^{t} p_j$ (such λ does exist), and $s_1 = s + \lambda f_n$. Then $f_n - s_1^2 = f_n - s^2 - \lambda f_n(2s + \lambda f_n)$, and $f_n - s_1^2 \notin p_1, \ldots, p_{r+l}$ by our choice of λ.

Now we claim that $f_n - s_1^2 \notin p_{r+l+1}, \ldots, p_t$. If $f_n - s_1^2 \in p_j$ for some $j \in \{r+l+1, \ldots, t\}$, then $2s + \lambda f_n \in p_j$. Notice that since B is a commutative ring containing a field k, either 2 is invertible in B or 2 is zero in B. If 2 is zero in B, then $\lambda f_n \in p_j$, which is impossible. If 2 is invertible in B, then $s + (1/2)\lambda f_n \in p_j$, which contradicts that $s - bf_n \notin p_k$ for $k > r$ and any $b \in I$. So the claim follows.

Therefor $f_n - s_1^2$ is a nonzero divisor in $B/(f_1, \dots, f_{n-1})$. By our choice of s_1, we have that $I = (f_1, \dots, f_n, s_1)$, $s_1(1 - s_1) \in (f_1, \dots, f_n)$, $s_1 \in I^2$, $I = (f_1, \dots, f_n) + I^2$, and $\{f_1, \dots, f_{n-1}, f_n - s_1^2\}$ is a regular sequence in I. $\square$

Throughout the rest of this paper, we will denote $P \otimes A[T]$ by $P[T]$ and use similar obvious notation.

Now we state our theorem for the case when dimension of $A \geq 4$, even. The proof of this theorem is motivated by [7 , Proposition 6.2, Mandal's online talk draft].

Theorem 3.4. *Let A be a commutative Noetherian ring of dimension n ($n \geq 4$, even) containing the field of rational numbers. Let I be a local complete intersection ideal of height n in $A[T]$. Suppose there is a surjection $\xi : P[T] \to I$, where P is a stably free A-module of rank n. Then I is a set theoretic complete intersection in $A[T]$.*

Proof. We give the proof of the theorem in several steps

Step 1. We want to find two local complete intersection ideals in $A[T]$ of height n, which are comaximal with I.

Since I/I^2 is free $A[T]/I$-module of rank n and I is a local complete intersection, we may write $I = (f_1, \dots, f_n) + I^2$, where $\{f_1, \dots, f_n\}$ is a regular sequence in $A[T]$. By lemma 3.3, we may assume the image of $f_n - s^2$ in $A[T]/(f_1, \dots, f_{n-1})$ is a nonzero divisor, $I = (f_1, \dots, f_n, s)$ and $s(1-s) \in (f_1, \dots, f_n)$. Let $K_1 = (f_1, \dots, f_n, 1 - s)$, then $K_1 \cap I = (f_1, \dots, f_n)$. Since $\{f_1, \dots, f_{n-1}, f_n - s^2\}$ is a regular sequence and $I = (f_1, \dots, f_{n-1}, f_n - s^2) + I^2$, we can write $(f_1, \dots, f_{n-1}, f_n - s^2) = I \cap K_2$ for some local complete intersection ideal K_2 in $A[T]$, which is comaximal with I. We may assume K_1, K_2 are ideals of height n, since if one of them equals $A[T]$, then I is a complete intersection. Let $g = f_n - s^2$, then $gA[T] + K_1 = A[T]$, and hence I, K_1, K_2 are pairwise comaximal. Let $I^{(2)} = (f_1, \dots, f_{n-1}) + I^2$, then $I^{(2)} \cap K_1 \cap K_2 = (f_1, \dots, f_{n-1}, gf_n)$. It is clear that $g = -s^2$ is unit modulo K_1 and $f_n = s^2$ is unit modulo K_2.

Step 2. We may assume ($f_1(0), \dots, f_{n-1}(0), g(0)f_n(0)$) is an ideal of height n in A.

To see this, we can replace T by $T+\lambda$ for some suitable $\lambda \in \mathbb{Q}$, and assume $(f_1(0),\ldots,f_{n-1}(0),g(0)f_n(0)) = A$ or $\mathrm{ht}(f_1(0),\ldots,g(0)f_n(0)) = n$. But if $(\ f_1(0),\ldots,f_{n-1}(0),g(0)f_n(0)\ ) = A$, then $I(0) = A$. Thus P has a umimodular element. It is then clear that I is a complete intersection in $A[T]$ by using lemma 2.3 as in the proof of theorem 3.2. Therefore we may assume $(\ f_1(0),\ldots,f_{n-1}(0),g(0)f_n(0)\ )$ is an ideal of hight n in A.

Step 3. We may assume $(f_1,\ldots,f_{n-1},gf_n)A(T)$ is an ideal of height n in $A(T)$.

If not, then $I^{(2)}$ contains a monic polynomial and hence so does I. From the surjection ξ, we see $P(T)$ has a unimodular element when localized at such a monic polynomial. It implies $P[T]$ has a unimodular element by [2, theorem 3.4]. Hence I is a complete intersection.

Step 4. We define a local orientation on $I^{(2)}$.

Since $(f_1,\ldots,f_{n-1},f_n) = I\cap K_1$ and K_1 is a local complete intersection ideal of height n and comaximal with I in $A[T]$, we have two natural surjective homomorphisms

$$\begin{array}{rcl} (A[T])^n & \overset{\alpha}{\to} & I\cap K_1 \\ (A[T])^n & \overset{\alpha'}{\to} & I\cap K_1 \end{array}$$

defined by $\alpha(e_i) = f_i$ for $i = 1,\ldots,n$, and $\alpha'(e_j) = f_j$ for $j = 1,\ldots,n-1$, $\alpha'(e_n) = -f_n$ where $e_1,\ldots,e_n$ is the standard basis of A^n . Since $(f_1,\ldots,f_{n-1},g) = I\cap K_2$ and $I^{(2)}\cap K_1\cap K_2 = (f_1,\ldots,f_{n-1},gf_n)$, where I,K_1,K_2 are pairwise comaximal, we also have two natural surjective homomorphisms

$$\begin{array}{rcl} (A[T])^n & \overset{\beta}{\to} & I\cap K_2 \\ (A[T])^n & \overset{\gamma}{\to} & I^{(2)}\cap K_1\cap K_2 \end{array}$$

defined by $\beta(e_i) = f_i$ for $i = 1,\ldots,n-1$, $\beta(e_n) = g$, and $\gamma(e_j) = f_j$ for $j = 1,\ldots,n-1$, $\gamma(e_n) = gf_n$. We define a local orientation ω of $I^{(2)}$ as $\omega = \gamma\otimes A[T]/I^{(2)}$.

Step 5. We want to show $\omega\otimes A[T]/T$ can be lifted to a surjection $\theta : A^n \to I^{(2)}(0)$.

Let $\xi(0) = \xi\otimes A[T]/T$, $\omega(0) = \omega\otimes A[T]/T$, $\omega_I(0) = \alpha\otimes A[T]/I\otimes A[T]/T$, $\omega_{K_1}(0) = \alpha\otimes A[T]/K_1\otimes A[T]/T$, and $\omega_{K_2}(0) = \beta\otimes A[T]/K_2\otimes A[T]/T$. We show $(I^{(2)}(0),\omega(0)) = 0$ in $E(A)$, the Euler class group of A. There are four cases:

Case 1. $K_1(0)$ and $K_2(0)$ are ideals of height n in A.

From the surjection α', we have the following relation in $E(A)$:

1. $(I(0), -\omega_I(0)) + (K_1(0), -\omega_{K_1}(0)) = 0$.

   From the surjection β, we have $(I(0), \omega_I(0)) + (K_2(0), \omega_{K_2}(0)) = 0$ in $E(A)$. Since $g(0) = -s^2(0)$ is unit modulo $K_1(0)$ and $f_n(0) = s^2(0)$ is unit modulo $K_2(0)$, from the surjections β and γ and lemma 5.4 in [3], we have the following two relations in $E(A)$:

2. $(I(0), \omega_I(0)) + (K_2(0), \omega_{K_2}(0)) = 0$, and

3. $(I^{(2)}(0), \omega(0)) + (K_1(0), -s^2(0)\omega_{K_1}(0)) + (K_2(0), s^2(0)\omega_{K_2}(0)) = (I^{(2)}(0), \omega(0)) + (K_1(0), -\omega_{K_1}(0)) + (K_2(0), \omega_{K_2}(0)) = 0$.

From the above three relations, we have $(I(0), \omega_I(0)) + (I(0), -\omega_I(0)) = (I^{(2)}(0), \omega(0))$ in $E(A)$.

Since $I(0)$ is the image of $\xi(0) : P \to I(0)$ and P is a stably free A-module, $(I(0), \omega_I(0)) + (I(0), -\omega_I(0)) = 0$ in $E(A)$ by [3, Proposition 6.2 and Corollary 7.9]. Thus $(I^{(2)}(0), \omega(0)) = 0$ in $E(A)$.

Case 2. $K_1(0) = A$ and $K_2(0)$ is an ideal of height n in A.

From the surjection α, we have $(I(0), \omega_I(0)) = 0$ in $E(A)$. Hence from the surjection β, we have $(K_2(0), \omega_{K_2}(0)) = 0$ in $E(A)$. Furthermore, the surjection γ implies $0 = (I^{(2)}(0), \omega(0)) + (K_2(0), s^2(0)\omega_{K_2}(0)) = (I^{(2)}(0), \omega(0)) + (K_2(0), \omega_{K_2}(0))$. So it is easy to see that $(I^{(2)}(0), \omega(0)) = 0$ in $E(A)$.

Case 3. $K_2(0) = A$ and $K_1(0)$ is an ideal of height n in A.

Similar proof as in case (2).

Case 4. $K_1(0) = A$ and $K_2(0) = A$.

In this case, $(I^{(2)}(0), \omega(0)) = 0$ in $E(A)$ by definition.

Therefore $\omega \otimes A[T]/T$ can be lifted to a surjection $\theta : A^n \to I^{(2)}(0)$ by theorem 2.7. And thus ω induces a surjection $\phi : A[T]^n \to I^{(2)}/{I^{(2)}}^2 T$ by [4, Remark 3.9].

Step 6. We now show ω can be lifted to a surjection $\psi : (A[T])^n \to I^{(2)}$.

By a similar argument as in step 5 and considering the Euler class group of $E(A(T))$ (Notice that $\dim A(T) = n$), it is easy to see $\omega \otimes A(T)$ can be lifted to a surjection $\phi' : A(T) \to I^{(2)}A(T)$. Thus by theorem 2.8, the surjection $\phi : A[T]^n \to I^{(2)}/{I^{(2)}}^2 T$ can be lifted to a surjection $\psi : A[T]^n \to I^{(2)}$.

Therefore, $I^{(2)}$ is a complete intersection, and thus I is a set theoretic complete intersection ideal in $A[T]$. The proof of the theorem is complete. □

Now we are able to give a proof of theorem 1.2 we stated in the Introduction in the case when n is bigger than 4 and even.

Theorem 3.5. *Let A be a commutative Noetherian ring of dimension n ($n \geq 4$, even) containing the field of rational numbers. Let I be a local complete intersection ideal of height n in $A[T]$, such that I/I^2 is free $A[T]/I$-module of rank n. Suppose $(A[T]/I)$ is torsion in $K_0(A[T])$. Then I is a set theoretic complete intersection in $A[T]$.*

Proof. By lemma 3.1, there exist a local complete intersection ideal K in $A[T]$ with height n, and a surjection $\xi : \tilde{P} \to K$ such that:

1. $K \subset I$, $\sqrt{K} = \sqrt{I}$,
2. $\tilde{P} \oplus A[T] \approx A[T]^{n+1}$.

By theorem 2.6. $\tilde{P}$ is extended from A, and thus $\tilde{P} \approx P[T]$, for some stably free A-module P of rank n. By theorem 3.4, K is a set theoretic complete intersection in $A[T]$. Therefore I is a set theoretic complete intersection in $A[T]$. The proof of the theorem is complete. □

Finally, let us state our theorem for the case when $\dim A=3$:

Theorem 3.6. *Let A be a commutative Noetherian ring of dimension 3 containing the field of rational numbers. Let I be a local complete intersection ideal of height 3 in $A[T]$, such that I/I^2 is free $A[T]/I$-module of rank 3. Suppose $(A[T]/I)$ is torsion in $K_0(A[T])$. Then I is a set theoretic complete intersection in $A[T]$.*

Proof. By lemma 3.1, there exist a local complete intersection ideal K in $A[T]$ with height 3, and a surjection $\xi : P \to K$ such that:

1. $K \subset I$, $\sqrt{K} = \sqrt{I}$,
2. $P \oplus A[T] \approx A[T]^4$.

From the surjection $\xi : P \to K$ and any given trivalization χ of $\bigwedge^3 P$, we get an element $(K, \omega_K) \in E(A[T])$ and $e(P, \chi) = (K, \omega_K)$ in $E(A[T])$ where (K, ω_K) is obtained from the pair (ξ, χ). By theorem 2.5, P has a unimodular element. Thus $e(P, \chi) = 0$ in $E(A[T])$ by [6, corollary 4.11]. It follows that $(K, \omega_K) = 0$ in $E(A[T])$, and thus K is generated by 3 elements by [6, theorem 4.7]. Therefore I is a set theoretic complete intersection in $A[T]$.

$\square$

Acknowledgement

I sincerely thank my advisor, Professor N. Mohan Kumar, for his guidance and many valuable suggestions, that have made this paper possible in its present form. I also wish to thank Dr. G.V. Ravindra for his encouragement.

References

[1] H. Bass, *Modules which support non-singular forms*, J. Algebra 13 (1969), 246–252.

[2] S.M. Bhatwadekar, R. Sriharan, *On a question of Roitman*, J. Ramanujan. Math. Society 16 (2001), 45–61.

[3] S.M. Bhatwadekar, R. Sriharan, *The Euler class group of a Noetherian ring*, Compositio Math. 122 (2000), 183–222.

[4] S.M. Bhatwadekar, R. Sriharan, *Projective generation of curves in polynomial extension of an affine domain and a question of Nori*, Invent. Math. 133 (1998), 161–1922.

[5] M. Boratynski, *A note on set-theoretic complete intersection ideals*, J. Algebra 54 (1978), 1–5.

[6] M. K. Das, *The Euler class group of a polynomial algebra*, J. Algebra 264 (2003), 582–612.

[7] S. Mandal, *Euler cycles, http://www.math.ku.edu/˜mandal/talks/talkEuler.pdf*, (2005)

[8] S. Mandal, *On set theoretic intersection in affine spaces*, J. Pure App. Algebra 51 (1988), 267–275.

[9] Mohan Kumar, N., *Complete intersections*, J. Math. Kyoto Univ. 17 (1977), 533–538.

[10] M.P. Murthy, *Zero cycles and projective modules*, Ann. Math. 140 (1994), 405–434.

[11] B. Plumstead, *The conjecture of Eisenbud and Evans*, Amer. J. Math. 105 (1983), 1417–1433.

[12] D. Quillen, *Projective modules over polynomial rings*, Invent. Math. 36(1976), 167–171.

[13] R. A. Rao, *The Bass-Quillen conjecture in dimension but characteristic $\neq$ 2, 3 via a question of A. Suslin*, Invent. Math. 93 (1988), 609–618.

[14] L. Szpiro, *Equations defining space curves*, Tata Institue, Bombay, 1979.